

Wednesday 17 June 2015 - Morning

A2 GCE HUMAN BIOLOGY

F225/01 Genetics, Control and Ageing

Candidates answer on the Question Paper.

OCR supplied materials:

None

Other materials required:

- Electronic calculator
- Ruler (cm/mm)

Duration: 2 hours



Candidate forename					Candidate surname				
Centre number						Candidate nu	umber		

INSTRUCTIONS TO CANDIDATES

- Write your name, centre number and candidate number in the boxes above. Please write clearly and in capital letters.
- Use black ink. HB pencil may be used for graphs and diagrams only.
- Answer all the questions.
- Read each question carefully. Make sure you know what you have to do before starting your answer.
- Write your answer to each question in the space provided. If additional space is required, you should use the lined pages at the end of this booklet. The question number(s) must be clearly shown.
- Do not write in the bar codes.

INFORMATION FOR CANDIDATES

- The number of marks is given in brackets [] at the end of each question or part question.
- The total number of marks for this paper is 100.
- Where you see this icon you will be awarded marks for the quality of written communication in your answer.
- You may use an electronic calculator.
- You are advised to show all the steps in any calculations.
- This document consists of 32 pages. Any blank pages are indicated.



2 BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

Answer **all** the questions.

Hor	meostasis has been defined as 'the maintenance of a <i>constant</i> or <i>stable</i> internal environment'.
(a)	Explain why the word 'stable' is more appropriate than the word 'constant' in a definition of homeostasis.
(b)	The autonomic nervous system plays an important role in homeostasis.
(b)	The box below contains some terms that could be used to describe the features or activity of the two branches of the autonomic nervous system.
	acetylcholine adrenaline decrease glycogenesis
	alvcogenolysis increase noradrenaline motor sensory

Complete Table 1.1 below by inserting the most appropriate term.

Terms may be used once, more than once or not at all.

Some boxes have been completed for you.

1

Factoria an activity	Branch of autonomic nervous system							
Feature or activity	Parasympathetic	Sympathetic						
Type of neurone	motor							
Process triggered in liver cells								
Effect on heart rate								
Neurotransmitter released at the SAN		noradrenaline						

[4]

(c) Control of body temperature is an important aspect of homeostasis. Conditions such as hypothermia can be fatal if not treated.

Fig. 1.1 compares the effects of three methods of treatment for mild hypothermia.




Fig. 1.1

(i)	Explain how shivering causes a rise in core body temperature.
	[3]
(ii)	Fig. 1.1 shows that placing a person with hypothermia in a bath of warm water results in an initial drop in core body temperature.
	Suggest why the core body temperature drops initially.
	[2]

(d) Re-warming by inhaling warmed air is also known as **core re-warming**. Fig. 1.2 shows the route taken by the warm air as it is inhaled by the person.




Fig. 1.2

Name the part of the brain in which the thermoregulatory centre is located.
[1]
Identify structure X.
[1]
Using the information in Fig. 1.2, suggest why core re-warming is better than other methods of re-warming at treating severe hypothermia.
In your answer, you should refer to the symptoms of severe hypothermia.
[3]

© OCR 2015 Turn over

[Total: 16]

- 2 Down's syndrome (DS) is a genetic condition that occurs in approximately 1 in every 800 births.
 - **Fig. 2.1a** shows a karyotype from a male with a rare form of DS. In this karyotype, there is an additional piece of genetic material attached to one copy of chromosome 14.

A larger diagram of both copies of chromosome 14 is shown in Fig. 2.1b.




Fig. 2.1b


		1
(a)	•	Describe how the appearance of the karyotype shown in Fig. 2.1a differs from that of the more common form of DS.
	•	Suggest when and how the transfer of extra genetic material to chromosome 14 occurred, and what events led to cells with this karyotype being formed.
		In your answer, you should include a comparison of the karyotypes and describe the events during and after meiosis that led to the formation of the karyotype in Fig. 2.1a.

.....[7]

(b) The National Cytogenic Register for DS was set up in the United Kingdom in 1989.

This register holds anonymous data on families for over 26 000 cases of DS. These cases were diagnosed either antenatally or postnatally.

Fig. 2.2 shows the changes in the number of DS cases diagnosed between 1990 and 2005. It also shows the changes in the number of live births of children with DS for the same period.



Key:

- total number of cases of DS diagnosed
- --- estimated number of live births assuming no antenatal screening or terminations were available
- actual number of live births when antenatal screening or terminations were available

Fig. 2.2

(i)	Suggest why the data on the National Cytogenic Register for DS are held anonymously.
	[1]
(ii)	Using the information in Fig. 2.2, suggest why the number of cases of DS diagnosed between 1990 and 2005 has changed.
	[2]
(iii)	Suggest what factor the investigators took into account in estimating the number of children with DS who would have been born , assuming that no antenatal screening or terminations were available.
	[1]
	[Total: 11]

3 GABA (gamma-aminobutyric acid) is one of the most common neurotransmitters in the human central nervous system.

GABA is synthesised from the amino acid glutamate.

(a) Fig. 3.1 is a diagram of GABA.




Fig. 3.1

GABA could also be described as an amino acid.

On Fig. 3.1, circle and name the **two** chemical groups on the GABA molecule which indicate that it is also an amino acid.

.....the answer must be drawn on Fig. 3.1......[1]

(b) GABA acts as an inhibitory neurotransmitter.

When GABA binds to its receptor, a chloride ion channel is opened and chloride ions (Cl^-) enter the neurone.

Fig. 3.2 represents two neurones forming synapses with a third neurone.




Fig. 3.2

Suggest how GABA is released from the inhibitory neurone **and**, using the information in Fig. 3.2, explain why an action potential may not occur in neurone **P** if both GABA and acetylcholine are released.

In your answer, you should make clear both the sequence of events in the release of the neurotransmitter and why an action potential could be prevented by GABA.
 [6]

(c)	GABA is the neurotransmitter in the parts of the brain responsible for sensations such as fear and anxiety. These sensations result from neurones becoming over-stimulated .
	Diazepam is a drug which is often prescribed to reduce anxiety. This drug can also be used to

reduce alcohol withdrawal symptoms and muscle spasms.

(i)	State two symptoms of alcohol withdrawal other than anxiety.
	1
	2[1]
(ii)	Name one other condition for which Diazepam could be used therapeutically.
	[1]
	[Total: 9]

13 BLANK PAGE

Question 4 begins on page 14

PLEASE DO NOT WRITE ON THIS PAGE

- 4 Some aspects of human physiology, behaviour and mental activity follow a pattern which cycles over a period of approximately 24 hours. This cyclical pattern is called the 'circadian rhythm'. It coincides approximately with periods of light and darkness.
 - (a) Circadian disorders, such as sleep-wake cycle disturbances, are associated with normal ageing.

What problems disturbances?	might	elderly	people	or their	carers	experience	as a	result of	sleep-wake
						•••••		•••••	•••••
									[3]

- (b) Circadian rhythm is regulated as follows:
 - a part of the brain called the SCN receives impulses from the optic nerve
 - the SCN regulates a gland in the brain which secretes a substance called **melatonin**
 - melatonin levels in blood are lowest during the day and highest at night
 - high levels of melatonin induce inactivity and sleep.

Fig. 4.1 is a summary of the interaction between the SCN and melatonin.




Fig. 4.1

(i)	What type of gland is W ?	
	What type of substance is melatonin?	
	Give a reason for your answers.	
		[2]
(ii)	Using the information in Fig. 4.1, suggest why circadian rhythm problems are more like to occur in the following groups of people:	cely
	those who are housebound	
	those with untreated glaucoma	
	those with Alzheimer's.	
		[3]

(c)	Stud	dies of melatonin levels in blood show large variations in concentrations between individuals.
	The	size of gland W has also been shown to vary between individuals.
	(i)	Suggest a technique that could be used to allow the size of gland ${\bf W}$ to be measured safely and repeatedly.
		Explain the reason for your suggestion.
		[2]
	(ii)	Longitudinal studies are studies of populations over time.
		Researchers have suggested that a longitudinal study would be a better experimental design for investigating the link between ageing, melatonin and changes in circadian rhythms.
		In such a study, identify the following:
		the independent variable
		a variable that would be controlled in a longitudinal study.
	/:::\	[2]
	(iii)	A study showed a negative correlation between melatonin concentration in blood and ageing.
		Complete the graph below to show this correlation.
		À
		———
		[2]

[Total: 14]

17 BLANK PAGE

Question 5 begins on page 18

PLEASE DO NOT WRITE ON THIS PAGE

The control of blood glucose is essential because changes in blood glucose concentration will change the water potential of blood.

	-	otential changes are detected by the osmoregulatory centre and adjustments are then the secretion of ADH.
(a)	(i)	State one reason, other than the effect that it has on water potential, why blood glucose concentrations must be controlled.
	(ii)	State the location in the brain of the osmoregulatory centre and the source of ADH secretion.
		osmoregulatory centre
		source of ADH secretion
		[1]
(b)	•	eople with type 1 diabetes do not administer the appropriate medication, the concentration lucose in the blood can remain higher than the normal range for relatively long periods of e.
	(i)	Name the medication that is required to treat type 1 diabetes.
		Describe how and when this medication is administered and how it acts to reduce the blood glucose concentration.
		[5]

5

(11)	You should assume that no glucose is lost from the blood and that glucose levels reinigh. Justify your prediction.	
	effect on ADH release	
	effect on urine volume	
	justification	
		 [31

Question 5(c) begins on page 20

(c) Fig. 5.1 represents the structure of part of a kidney nephron and its surrounding blood vessels.




Fig. 5.1

(i)	Using Fig. 5.1, identify, with a reason, the following:
	the region of the nephron shown
	reason
	the type of blood vessel shown
	reason
	[2]
(ii)	Structure T is a 'tight junction' between two adjacent cells. Tight junctions prevent proteins located at the surface of cells from changing their positions.
	Suggest why, if there were no tight junctions, proteins in cell surface membranes would be able to change positions.
	[2]

(iii)	The region of the nephron shown in Fig. 5.1 is the region where glucose is reabsorbed.
	Using the information in Fig. 5.1, name a mechanism used for selective reabsorption of glucose and explain why glucose appears in urine when blood glucose concentrations are high.
	[3]
` '	me people with diabetes will develop a condition called diabetic kidney disease. One of the nptoms of this condition is the appearance of the protein albumin in the urine.
(i)	Suggest where albumin is normally found in the body.
	[1]
(ii)	Suggest, with reasons, which region of the kidney is damaged due to diabetic kidney disease.
	[2]
	[2] [Total: 20]

22 BLANK PAGE

PLEASE DO NOT WRITE ON THIS PAGE

- 6 The age at which a woman goes through the menopause depends on several factors.
 - Smoking, unemployment and being at higher risk of coronary heart disease are factors linked to a lower age of menopause.
 - Having children, use of oral contraception and having a lower BMI are factors linked to a higher age of menopause.
 - The age of menopause varies between different ethnic groups.

In a study involving large populations of women, the median age of 'natural menopause' in a population was found to be 50.4 years.

(a)	(i)	Suggest what is meant by natural menopause.
		[2]
	(ii)	A median value for a large population is obtained as follows:
		all the values obtained are placed in order from lowest to highest
		the median is the middle value in the range.
		Suggest why the mean age of menopause can vary significantly between populations but the median does not.
		[2]
(b)		emicals in cigarette smoke have been shown to cause damage to cells in ovaries. Damaged s are no longer able to perform their normal function.
	Exp	plain how damage to cells in the ovaries could result in an earlier onset of the menopause.
		[7]

(c) Women may become infertile several years before the menopause. Tests which predict the timing of the menopause would allow women to plan their pregnancies.

Recent genetic studies have identified *loci* which are associated with variation in the age of onset of natural menopause.

- Each locus is a short DNA sequence which differs by one nucleotide
- One variant increases the risk of an early menopause

Fig. 6.1 is a diagram of a DNA sequence

- The variation occurs at the point indicated by the arrow where the nucleotide can be either 'C' or 'T'
- The presence of 'C' at this point increases the risk of an early menopause.



	region Z
	Fig. 6.1
(i)	What do the letters C and T represent in Fig. 6.1?
	c
	Т[1]
(ii)	Region Z is a nucleotide sequence that is complementary to the active site of an enzyme known as EcoRI. EcoRI is a type of enzyme that will cut the DNA strand.
	State the following:
	the type of enzyme that cuts DNA at a particular recognition site
	the nature of the nucleotide sequences at the recognition site
	the name of the bond in the DNA strand that the enzyme breaks
	[3]

(iii) Separate DNA samples of the 'C' variant and the 'T' variant were treated using EcoRI. The DNA was analysed using electrophoresis.

Describe **and** explain how the appearance of the pattern on an electrophoresis gel will differ between the DNA samples.


You should assume that:

region Z occurs only once
no other recognition sites for EcoRI are present.
[3]
[Total: 13]

Question 7 begins on page 26

7 The appearance of blood samples viewed under a microscope can be used to diagnose a number of different conditions.

Fig. 7.1 shows a blood sample. An abnormal cell known as a **schistocyte** is labelled. Schistocytes are formed from fragments of red blood cells.



magnification × 2800

Fig. 7.1

(a) (i) Use the information in Fig. 7.1 to calculate the size of cell A.Show your working. Give your answer to the nearest whole number.

Answer = μm [2]

	(ii)	When activated platelets bind to the endothelium of small arterioles and capillaries, schistocytes are formed.
		Suggest why the binding of activated platelets leads to the formation of schistocytes.
		[1]
	(iii)	State one genetic disease that could be diagnosed by examination of a blood smear.
		[1]
(b)		nfection by a strain of the bacterium <i>E. coli</i> called O157 can lead to the appearance of stocytes in a blood sample.
		ctions by this bacterium are the most common cause of acute kidney failure in children. results in a condition called HUS .
		t children who develop HUS make a full recovery. A few children are left with permanent ey damage.
	(i)	Name the short-term treatment given to children with acute kidney failure and outline the alternative long-term treatment options that would be available if necessary.
		rea

Question 7(b)(ii) begins on page 28

	(11)	anaemic. Explain your suggestion.
		[3]
(c)		ne forms of HUS are not associated with <i>E. coli</i> O157 infections. These are known as pical HUS or AHUS .
	Son	ne cases of AHUS are known to be due to an inherited gene mutation.
		efully read and then complete the following passage about the inheritance of AHUS and role of the genetic counsellor.
	Som	ne inherited forms of AHUS are due to a mutant allele.
	Pare	ents show no symptoms of the disease so, if they have a child who develops AHUS,
	the	child must be for this allele. Since neither parent
	shov	ws symptoms, yet both carry the mutant allele, the allele does not show
		linkage.
		enetic counsellor would use a diagram to explain the risk
		ny further children either developing AHUS or being a carrier. Where unaffected parents
		e one child with AHUS , there would be a
	OI II	aving a child who is a carrier. [5]

[Total: 17]

END OF QUESTION PAPER

ADDITIONAL ANSWER SPACE

If additional answer space is required, you should use the following lined page(s). The question number(s) must be clearly shown in the margins.			

•••••	
•••••	

•••••	
•••••	



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.